

Assessing the Allergic Potential of Indoor Air Fungal Contaminants



Part of a Cross-Laboratory Study: Children's Health Problems Associated with Fungal Exposure

Marsha D W Ward, Michael E Viana², Yongjoo Chung³, Najwa Haykal-Coates, Lisa B Copeland, Stephen H Gavett, and MaryJane K Selgrade

²NCSU, CVM, Raleigh, NC, ³ UNC, SPH, Chapel Hill, NC

INTRODUCTION

- **ENVIRONMENTAL ISSUE:** Dampness in indoor environments contributes to fungal growth in both home and workplace buildings.
- Fungi are the principal microorganisms responsible for the health problems in such buildings.
- Children spend >90% of their time indoors including home, school,
- Indoor exposures of children to molds results in complex and controversial health effects.
- The toxigenic fungi Stachybotrys chartarum was chosen as prototype mold for these studies.

SCIENTIFIC APPROACH:

The cross-laboratory approach provides a multi-faceted assessment and evaluation of exposure and health effects of exposure to indoor molds by using a combination of analytical, experimental, epidemiological, and modeling approaches.

PARTICIPATING LABORATORIES AND CENTERS

NATIONAL EXPOSURE RESEARCH LABORATORY

NATIONAL HEALTH AND ENVIRONMENTAL **EFFECTS RESEARCH LABORATORY**

Experimental Toxicology Division

- Establish a mouse model of allergic lung disease to S. chartarum Identify, characterize and compare IgE inducing proteins from three fungi (Metarhizium anisopliae, S. chartarum, and Penicillium
- Demonstrate parallels between human and rodent responses

NATIONAL RISK MANAGEMENT RESEARCH **LABORATORY**

NATIONAL CENTER FOR ENVIRONMENTAL **ASSESSMENT**





S. chartarum spores


Fifty day-old BALB/c mice (Charles River Laboratories,

METHODS

Fungus was ground with a mortar and pestle, stirred overnight, clarified by centrifugation, sterile filtered, and concentrated to yield a crude extract:

- MACA equal protein amounts of *M. anisopliae* mycelial, conidial, inducible protease/chitinase extracts
- SCE-1 a mixture of 5 S. chartarum isolates Mice received one of the following treatments by involuntary aspiration (IA) on the schedule described:
- 4 doses of MACA or SCE-1 in HBSS
- 4 doses of HBSS (vehicle control)
- 4 doses of BSA in HBSS (negative control) 3 doses of HBSS and 1 dose of MACA or SCE-1
- (inflammatory control)


Mice were killed on day 1 and day 3 by an overdose of sodium pentobarbital and exsanguinated by cardiac





- Respiratory Physiology
- Immediate respiratory physiological responses were measured after each aspiration (D-28,D-14,D-7,D0) by barometric whole-body plethysmography (Buxco
- Airway hyperresponsiveness was measured on day 1 and 3 by exposure to doubling concentrations of nebulized methacholine (Buxco system).
- Antibody and cytokine ELISAs
- Antibody and cytokine levels were quantified by ELISA.
- Lungs were fixed in 10% buffered formalin acetate. embedded in paraffin, and sectioned. Sections were examined and scored by a board-certified veterinary pathologist (Veritas Laboratories, Inc., Burlington, NC).
- Data were analyzed for statistical significance with SAS (SAS Corporation, Cary, NC), using a one-way analysis of

RESULTS


Representative examples from *Stachybotyrs chartarum* (SCE-1) and Metarhizium anisopliae (MACA) studies are presented









Multiple Exposures to Fungal Extract Increased BALF IL-5 and Serum Total IgE




Multiple Exposures to Fungal Extract **Increased Lung Influx of Inflammatory and Lymphocytic Cells**



Multiple Exposures to MACA Resulted in **Increased Immediate Respiratory Responses and Airway Hyperresponsiveness**



Several IgE Inducing Proteins Have Been Identified in MACA and SCE-1 (not shown)



proteins in MACA are allergens.

CONCLUSION

- Repeated pulmonary exposures to either Stachybotrys chartarum (SCE-1) and *Metarhizium anisopliae* (MACA) Extracts Induced:
- Non-specific lung injury
- Permeability BALF total protein
- Cell damage BALF LDH activity Neutrophilic infiltration
- Increased antigen specific immune responses
- Increased BALF IL-5 and eosinophilic infiltration
- Serum total IqE
- Immediate airway reactivity (extract exposure) and hyperresponsiveness to non-specific challenge (methacholine)
- Both of these fungal extracts have demonstrated the ability to induce responses in the mouse characteristic of allergic lung disease in
- The mouse model provides the means to
- Identify and characterize the specific allergens
- Assess relative potency compared to other indoor allergens

IMPACT

The data generated

- Establishes methodology leading to risk assessment for children/infants from exposure to S. chartarum specifically but serves as a model for assessment of other indoor biological contaminants.
- Provides comprehensive information to be used by the Office of Air and Radiation and by the Regional Offices in helping our clients, the American parent, on dealing with this major environmental problem.

FUTURE DIRECTIONS

- Continue identification and characterization of IgE inducing proteins from three fungi in collaboration with NERL scientists.
- Demonstrate parallels between human and rodent responses
- Determine relative potency of the IgE inducing proteins compared
- Among the fungal "allergens"
- To better characterized allergens, i.e. house dust mite, cockroach, alcalase (benchmarked detergent enzyme)

ACKNOWLEDGEMENTS

US EPA

Denise M. Sailstad Elizabeth Boykin Judy Richards Debora L. Andrews Christopher A. Manuel Jennifer Vu

Donald L. Doerfler

